

PRODUCT MONOGRAPH

^{Pr}Vinorelbine Injection, USP

Sterile Solution for Injection

10 mg vinorelbine (as vinorelbine tartrate) per mL

Solution Must Be Diluted Before Use

Antineoplastic Agent

Fresenius Kabi Canada Ltd.

165 Galaxy Blvd, Suite 100

Toronto, ON M9W 0C8

Date of Revision:

August 14, 2023

Submission Control No: 273007

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	4
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	6
DRUG INTERACTIONS	12
DOSAGE AND ADMINISTRATION.....	12
OVERDOSAGE	14
ACTION AND CLINICAL PHARMACOLOGY	15
STORAGE AND STABILITY.....	16
SPECIAL HANDLING INSTRUCTIONS	16
DOSAGE FORMS, COMPOSITION AND PACKAGING	17
PART II: SCIENTIFIC INFORMATION.....	18
PHARMACEUTICAL INFORMATION.....	18
CLINICAL TRIALS.....	18
DETAILED PHARMACOLOGY	20
TOXICOLOGY	23
REFERENCES	30
PART III: CONSUMER INFORMATION.....	33

PrVinorelbine Injection, USP

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form/Strength	All Nonmedicinal Ingredients
Intravenous	Solution / 10 mg vinorelbine (as vinorelbine tartrate) per mL	Water for Injection

INDICATIONS AND CLINICAL USE

Vinorelbine Injection, USP (vinorelbine tartrate) is indicated for:

- treatment of advanced non-small cell lung cancer (NSCLC), as a single agent or in combination.
- treatment of patients with metastatic breast cancer who have failed standard first-line chemotherapy for metastatic disease.
- treatment of patients with metastatic breast cancer who have relapsed within 6 months of anthracycline-based adjuvant therapy.

Geriatrics (≥ 65 years of age): No overall differences in effectiveness or safety were observed between patients 65 years of age or greater and younger patients. Other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out (see WARNINGS AND PRECAUTIONS).

Pediatrics: Safety and effectiveness in children have not been established.

CAUTION:

VINORELBINE INJECTION, USP (vinorelbine tartrate) IS CYTOTOXIC DRUGS AND SHOULD BE USED ONLY BY PHYSICIANS EXPERIENCED WITH CANCER CHEMOTHERAPEUTIC DRUGS. BLOOD COUNTS SHOULD BE TAKEN PRIOR TO EACH DOSE. THE DOSAGE SHOULD BE REDUCED OR THE DRUG DISCONTINUED UPON EVIDENCE OF ABNORMAL DEPRESSION OF THE BONE MARROW.

This preparation is for intravenous administration only. Intrathecal administration of other vinca alkaloids has resulted in death. Syringes containing this product should be labelled "WARNING - FOR INTRAVENOUS USE ONLY. FATAL if given intrathecally."

CONTRAINDICATIONS

- Vinorelbine Injection, USP (vinorelbine tartrate) is contraindicated in patients with known hypersensitivity to vinorelbine tartrate.
- As with other vinca alkaloids, Vinorelbine Injection, USP is contraindicated in patients who have drug-induced severe granulocytopenia or severe thrombocytopenia.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

Vinorelbine Injection, USP (vinorelbine tartrate) is cytotoxic drugs and should be used only by physicians experienced with cancer chemotherapeutic drugs. Blood counts should be taken prior to the next dose. Discontinue or reduce the dosage upon evidence of abnormal depression of the bone marrow.

General

Vinorelbine Injection, USP is for intravenous use only. Vinorelbine tartrate is a moderate vesicant and can produce phlebitis or extravasation injury. Inadequate flushing of the vein after peripheral administration may increase the risk of phlebitis.

It is extremely important that the needle be properly positioned in the vein before this product is injected. If leakage into surrounding tissue should occur during intravenous administration of Vinorelbine Injection, USP, it may cause severe irritation. The injection should be discontinued immediately, and any remaining portion of the dose should then be introduced into another vein.

A low incidence of death (1%) due to neutropenic sepsis has been reported (see ADVERSE REACTIONS). Bone marrow toxicity, specifically granulocytopenia, is dose-limiting. Complete blood counts with differentials should be performed and results reviewed prior to each dose of Vinorelbine Injection, USP. Vinorelbine Injection, USP should not be administered to patients with granulocyte counts < 1000 cells/mm³. Patients developing severe granulocytopenia should be monitored carefully for evidence of infection and/or fever (see DOSAGE AND ADMINISTRATION).

In all instances where the use of Vinorelbine Injection, USP (vinorelbine tartrate) is considered for chemotherapy, the physician must evaluate the need and usefulness of the drug against the risk of adverse events. Most drug-related adverse reactions are reversible. If severe adverse events occur, the drug should be reduced in dosage or discontinued and appropriate corrective measures should be taken based on the clinical judgment of the physician. Reinstitution of therapy with Vinorelbine Injection, USP should be carried out with caution and alertness as to possible recurrence of toxicity.

Patients should be informed that Vinorelbine Injection, USP is vesicant and can produce phlebitis or extravasation injury, and that the major acute toxicities of vinorelbine are related to bone marrow toxicity, specifically granulocytopenia with increased susceptibility to infection, and neuropathy. They should also be advised to report fever or chills immediately. Vinorelbine

Injection, USP should not be used in pregnancy unless the physician feels the potential benefit justifies the risk of potential harm to the fetus.

Hematologic

Vinorelbine Injection, USP should be used with extreme caution in patients whose bone marrow reserve may have been compromised by prior irradiation or chemotherapy, or whose marrow function is recovering from previous chemotherapy.

Since dose-limiting clinical toxicity is the result of depression of the white blood cell count, it is imperative that complete blood counts with differentials be obtained prior to each dose of Vinorelbine Injection, USP (See ADVERSE REACTIONS, Hematologic).

Hepatic

There is no evidence that the toxicity of vinorelbine tartrate is enhanced in patients with elevated liver enzymes; no data are available for patients with severe baseline cholestasis. However, pharmacologic evidence suggests that the liver plays an important role in the metabolism of vinorelbine tartrate. Although there are no data available from patients with severe liver disease, caution should be exercised when administering Vinorelbine Injection, USP to patients with severe hepatic injury or impairment.

Neurologic

Patients with a prior history or pre-existing neuropathy, regardless of etiology, should be monitored for new or worsening signs and symptoms of neuropathy while receiving Vinorelbine Injection, USP.

Ophthalmologic

Care must be exercised to avoid contamination of the eye with Vinorelbine Injection, USP. Accidental exposure should be treated immediately with a large volume of irrigation solution (water or sodium chloride).

Radiation

Administration of Vinorelbine Injection, USP to patients with prior radiation therapy may result in radiation recall reactions (see ADVERSE REACTIONS and DRUG INTERACTIONS).

Respiratory

Acute shortness of breath and severe bronchospasm have been reported infrequently following the administration of vinorelbine tartrate and of other vinca alkaloids. These events have been encountered most commonly when the vinca alkaloid was used in combination with mitomycin and may require aggressive treatment, particularly when there is pre-existing pulmonary dysfunction. Bronchodilators, steroids and/or oxygen have produced symptomatic relief.

Sensitivity

Vinorelbine Injection, USP should be used with extreme caution in patients whose bone marrow reserve may have been compromised by prior irradiation or chemotherapy, or whose marrow function is recovering from previous chemotherapy.

Administration of Vinorelbine Injection, USP to patients with prior radiation therapy may result in radiation recall reactions (see ADVERSE REACTIONS and DRUG INTERACTIONS).

Special Populations

Pregnant Women: There are no studies in pregnant women. Vinorelbine has been shown to be embryotoxic and/or fetotoxic in animals (see TOXICOLOGY). Vinorelbine Injection, USP should not be used in pregnancy.

Nursing Women: It is not known whether the drug is excreted in human milk. Because many drugs are excreted in human milk and because of its potential for serious adverse reactions in nursing infants, it is recommended that nursing be discontinued in women who are receiving therapy with Vinorelbine Injection, USP.

Pediatrics: Safety and effectiveness in children have not been established.

Geriatrics: Of the total number of patients in North American clinical studies of intravenous vinorelbine tartrate, approximately one-third were 65 years of age or greater. No overall differences in effectiveness or safety were observed between these patients and younger patients. Other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Data in the following tables are based on the experience of 365 patients (143 patients with NSCLC; 222 patients with advanced breast cancer) for whom a complete safety database was available and who were treated with vinorelbine tartrate as a single agent in three North American trials (one NSCLC trial and two advanced breast cancer trials). Patients treated for breast cancer were allowed to have received adjuvant chemotherapy in both trials, and in one, up to two prior regimens for advanced disease. The dosing schedule was 30 mg/m² intravenous vinorelbine tartrate on a weekly basis.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug

reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Table 1: Summary of Adverse Events Occurring in $\geq 5\%$ of 365 Patients Receiving Single-Agent Vinorelbine Tartrate that are Possibly Attributable to the Study Medication^{a,b}

Adverse Event	% Incidence All Grades		% Incidence Grade 3		% Incidence Grade 4	
	ABC n = 222	NSCLC n = 143	ABC n = 222	NSCLC n = 143	ABC n = 222	NSCLC n = 143
General						
Injection site reaction	21	38	1	5	0	0
Asthenia	41	25	8	5	0	0
Pain	16	15	3	2	0	0
Pain at injection site	18	13	3	1	0	0
Fever	19	10	1	0	0	0
Pain (Abdomen)	12	6	1	1	0	0
Pain (Chest)	8	5	1	2	0	0
Phlebitis	5	10	0	1	0	0
Digestive System						
Nausea	50	33	3	1	0	0
Constipation	38	28	3	2	0	0
Anorexia	19	16	< 1	2	0	0
Stomatitis	16	15	0	0	0	0
Vomiting	23	14	2	1	0	0
Diarrhea	20	13	< 1	1	0	0
Musculoskeletal System						
Myasthenia	9	5	2	1	< 1	0
Nervous System						
Paresthesia	20	11	0	1	0	0
Hypoesthesia	11	10	< 1	0	< 1	0
Respiratory System						
Dyspnea	9	3	1	2	1	0

Adverse Event	% Incidence All Grades		% Incidence Grade 3		% Incidence Grade 4	
	ABC	NSCLC	ABC	NSCLC	ABC	NSCLC
	n = 222	n = 143	n = 222	n = 143	n = 222	n = 143
Skin and Appendages						
Alopecia	12	12	0	1	0	0
Rash	5	5	0	0	0	0

ABC = Advanced Breast Cancer; NSCLC = Non-Small Cell Lung Cancer

^a Grade based on modified criteria of the National Cancer Institute

^b Patients with NSCLC had not received prior chemotherapy. The majority of patients with advanced breast cancer had received prior chemotherapy.

Cardiovascular

Chest pain was reported in 5% of NSCLC and 8% of breast cancer patients. Most reports of chest pain were in patients who had either a history of cardiovascular disease or tumour within the chest. There have been rare reports of myocardial infarction; however, these have not been shown definitely attributable to vinorelbine tartrate.

Dermatologic

Alopecia was reported in only 12% of patients and was usually mild.

Vinorelbine tartrate is a moderate vesicant. Injection site reactions, including erythema, pain at injection site, and vein discoloration occurred in approximately one-third of all patients; 2% were severe. Chemical phlebitis along the vein proximal to the site of injection was reported.

Gastrointestinal

Mild or moderate nausea occurred in 32% of NSCLC and 47% of breast cancer patients treated with vinorelbine tartrate. Severe nausea was infrequent (1% and 3% in NSCLC and breast cancer patients respectively). Prophylactic administration of antiemetics was not routine in patients treated with single-agent vinorelbine tartrate. Constipation occurred in approximately 28% of NSCLC and 38% of breast cancer patients, with paralytic ileus occurring in less than 2% of patients. Vomiting, diarrhea, anorexia, and stomatitis were usually mild or moderate and occurred in less than 20% of patients.

Hepatic

Transient elevations of liver enzymes were reported without clinical symptoms.

Neurologic

Mild to moderate peripheral neuropathy manifested by paresthesia and hypoesthesia were the most frequently reported neurologic toxicities (10% to 20%, see Table 2). Loss of deep tendon reflexes occurred in less than 5% of patients. The development of severe peripheral neuropathy was unusual.

Pulmonary

Shortness of breath was reported in 3% of NSCLC and 9% of breast cancer patients; and was severe in 2% of each patient population. Interstitial pulmonary changes have been documented in a few patients.

Other

Asthenia occurred in approximately 25% of patients with NSCLC and 41% of patients with breast cancer. It was usually mild or moderate but tended to increase with cumulative dosing.

Other toxicities that have been reported in $\leq 5\%$ of patients include jaw pain, myalgia, arthralgia, headache, dysphagia, and rash. Hemorrhagic cystitis and the syndrome of inappropriate ADH secretion were each reported in 4% of patients. The treatment of these entities are mainly symptomatic. The treatment of hemorrhagic cystitis is intravenous fluids for forced diuresis and/or irrigation of bladder. For the treatment of SIADH, please refer to the major textbooks of medicine.

Abnormal Hematologic and Clinical Chemistry Findings

Hematologic

Granulocytopenia was the major dose-limiting toxicity with vinorelbine tartrate; it was generally reversible and not cumulative over time. Granulocyte nadirs occurred 7 to 10 days after the dose and usually recovered within the following 7 to 14 days. Granulocytopenia resulted in hospitalizations for fever and/or sepsis in 8% of NSCLC and 9% of breast cancer patients. Septic deaths occurred in approximately 1% of patients.

Grade 3 or 4 anemia occurred in 1% of lung cancer and 14% of breast cancer patients. Blood products were administered to 18% of patients who received vinorelbine tartrate. The incidence of Grade 3 and 4 thrombocytopenia was less than 1%.

Table 2: Hematologic Adverse Events and Clinical Chemistry Elevations in 365 Patients Receiving Single-Agent Vinorelbine Tartrate that are Possibly Attributable to the Study Medication^{a,b}

Hematology		ABC (%)	NSCLC (%)
Granulocytopenia	< 2000 cells/mm ³	96	80
	< 500 cells/mm ³	41	28

Hematology		ABC (%)	NSCLC (%)
Leukopenia	< 4000 cells/mm ³	99	81
	< 1000 cells/mm ³	16	12
Thrombocytopenia	< 100 000 cells/mm ³	6	4
	< 50 000 cells/mm ³	< 1	1
Anemia	< 11 g/dL Hgb	87	77
	< 8 g/dL Hgb	14	1
Hospitalizations due to granulocytopenic complications		9	8

Chemical Chemistry Elevations	% Incidence All Grades		% Incidence Grade 3		% Incidence Grade 4	
	ABC	NSCLC	ABC	NSCLC	ABC	NSCLC
Total Bilirubin						
NSCLC: n = 137	14	9	4	3	3	2
ABC: n = 214						
SGOT						
NSCLC: n = 133	74	54	7	2	< 1	1
ABC: n = 213						

ABC = Advanced Breast Cancer; NSCLC = Non-Small Cell Lung Cancer

^a Grade based on modified criteria of the National Cancer Institute

^b Patients with NSCLC had not received prior chemotherapy. The majority of patients with advanced breast cancer had received prior chemotherapy.

Observed During Clinical Practice

In a randomized study in NSCLC patients, 206 patients received treatment with vinorelbine tartrate plus cisplatin and 206 patients received single-agent vinorelbine tartrate. The incidence of severe nausea and vomiting was 30% for vinorelbine tartrate /cisplatin compared to < 2% for single-agent vinorelbine tartrate. Cisplatin did not appear to increase the incidence of neurotoxicity observed with single-agent vinorelbine tartrate. However, myelosuppression, specifically Grade 3 and 4 granulocytopenia, was greater with the combination of vinorelbine tartrate /cisplatin (79%) than with single-agent vinorelbine tartrate (53%). The incidence of fever and infection may be increased with the combination.

Post-Market Adverse Drug Reactions

In addition to adverse events reported from clinical trials, the following events have been identified during post-marketing use of vinorelbine tartrate. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. These events have been chosen for inclusion due to combination of their seriousness, frequency of reporting, or potential causal connection to vinorelbine tartrate or a combination of these factors.

Body as a Whole: Systemic allergic reactions reported as anaphylaxis, pruritis, urticaria and angioedema, flushing and radiation recall events such as dermatitis and esophagitis (see WARNINGS AND PRECAUTIONS) have been reported.

Cardiovascular: Hypertension, hypotension, vasodilation, tachycardia, and pulmonary edema have been reported.

Gastrointestinal: Dysphagia and mucositis have been reported.

Hematologic: Thromboembolic events including pulmonary embolus and deep venous thrombosis have been reported primarily in seriously ill and debilitated patients with known predisposing risk factors for these events.

Musculoskeletal: Headache has been reported with and without other musculoskeletal aches and pains.

Neurologic: Peripheral neurotoxicities such as, but not limited to, muscle weakness and disturbance of gait have been observed in patients with and without prior symptoms. Vestibular and auditory deficits have been observed with vinorelbine tartrate, usually when used in combination with cisplatin. There may be increased potential for neurotoxicity in patients with pre-existing neuropathy, regardless of etiology, who receive vinorelbine tartrate. Patients who receive vinorelbine tartrate and paclitaxel, either concomitantly or sequentially, should be monitored for signs and symptoms of neuropathy (see WARNINGS AND PRECAUTIONS).

Pulmonary: Pneumonia has been reported.

Vinorelbine can produce acute and subacute pulmonary reactions. The acute reaction usually resembles an allergic event and may respond to bronchodilators. Subacute pulmonary reactions occur shortly after drug administration and may be characterized by cough, dyspnea, hypoxemia, and interstitial infiltration. Subacute pulmonary reactions may respond to corticosteroid therapy.

Skin: Injection site reactions, including localised rash and urticaria, blister formation and skin sloughing have been observed in clinical practice. Some of these reactions may be delayed in appearance.

Other: Pain in tumour-containing tissue, back pain and abdominal pain have been reported. Electrolyte abnormalities including hyponatremia consistent with the syndrome of inappropriate ADH secretion, have been reported in seriously ill and debilitated patients.

Combination Use: Patients with prior exposure to paclitaxel and who have demonstrated neuropathy should be monitored closely for new or worsening neuropathy. Patients who have experienced neuropathy with previous drug regimens should be monitored for symptoms of neuropathy while receiving vinorelbine tartrate. Vinorelbine tartrate may result in radiosensitising effects with prior or concomitant radiation therapy (see WARNINGS AND PRECAUTIONS).

DRUG INTERACTIONS

Drug-Drug Interactions

Acute pulmonary reactions have been reported with vinorelbine tartrate and other vinca alkaloids used in conjunction with mitomycin (see WARNINGS AND PRECAUTIONS, General).

Vinorelbine Injection, USP should be administered with caution in combination with mitomycin. Although the pharmacokinetics of vinorelbine are not influenced by the concurrent administration of cisplatin, the incidence of toxicities, specifically granulocytopenia, high frequency hearing loss and tinnitus, with the combination of vinorelbine tartrate and cisplatin are higher than with single-agent vinorelbine tartrate.

Patients who receive vinorelbine tartrate and paclitaxel, either concomitantly or sequentially, should be monitored for signs and symptoms of neuropathy.

Administration of Vinorelbine Injection, USP to patients with prior or concomitant radiation therapy may result in radiosensitising effects.

Drug-Food Interactions

Interactions with food have not been established.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Interactions with laboratory tests have not been established.

DOSAGE AND ADMINISTRATION

Dosing Considerations

This preparation is for intravenous administration only. It should be administered by individuals experienced in the administration of cancer chemotherapeutic drugs.

Recommended Dose and Dosage Adjustment

The usual initial dose of Vinorelbine Injection, USP (vinorelbine tartrate) is 30 mg/m² administered weekly. The recommended method of administration is an intravenous injection over 6 to 10 minutes. In controlled trials, single-agent vinorelbine tartrate was given weekly until progression or dose-limiting toxicity.

No dose adjustments are required for renal insufficiency. If moderate or severe neurotoxicity develops, Vinorelbine Injection, USP should be discontinued. The dosage should be adjusted according to hematologic toxicity or hepatic insufficiency.

Dose Modifications for Hematologic Toxicity: Granulocyte counts should be ≥ 1000 cells/mm³ prior to the administration of Vinorelbine Injection, USP. In the referenced North American trial, in which hematologic adverse events were observed, the following dose adjustment scheme was employed and should be followed in patients receiving vinorelbine tartrate.

Table 3: Dose Adjustments Based on Granulocyte Counts

Granulocytes (cells/mm³) on Days of Treatment	Dose of Vinorelbine tartrate (mg/m²)
≥ 1500	30
1000 to 1499	15
< 1000	Do not administer. Repeat granulocyte count in 1 week. If granulocyte count is < 1000 cells/mm ³ for > 3 weeks, discontinue Vinorelbine Injection, USP.
Note: For patients who, during treatment with vinorelbine tartrate, have experienced fever and/or sepsis while granulocytopenic or required a delay in dosing of up to 3 weeks due to granulocytopenia, the dose of vinorelbine should be: 22.5 mg/m ² for granulocytes ≥ 1500 cells/mm ³ 11.25 mg/m ² for granulocytes 1000 to 1499 cells/mm ³	

Vinorelbine Injection, USP should be administered with caution to patients with hepatic insufficiency. In patients who develop hyperbilirubinemia during treatment with vinorelbine, the dose should be adjusted for total bilirubin.

Administration

Vinorelbine Injection, USP must be administered intravenously. It is extremely important that the intravenous needle or catheter be properly positioned before any vinorelbine is injected. Leakage into surrounding tissue during intravenous administration of vinorelbine may cause considerable irritation, local tissue necrosis and/or thrombophlebitis. If extravasation occurs, the injection should be discontinued immediately, and any remaining portion of the dose should then be introduced into another vein. The application of moderate heat to the area of leakage in the form of warm compress applied for 15 to 20 minutes at least four times per day for the first 24 to 48 hours in addition to rest and elevation of the affected site for 48-72 hours has been reported to help disperse drug and minimize discomfort associated with the extravasation of other vinca alkaloids.

As with other toxic compounds, caution should be exercised in handling and preparing the solution of Vinorelbine Injection, USP. Skin reactions may occur with accidental exposure. The use of gloves is recommended. If the solution of Vinorelbine Injection, USP contacts the skin or mucosa, immediately wash the skin or mucosa thoroughly with soap and water. Severe irritation of the eye has been reported with accidental contamination of the eye with another vinca

alkaloid. If this happens with Vinorelbine Injection, USP, the eye should be washed with water immediately and thoroughly.

Preparation for Administration:

Vinorelbine Injection, USP must be diluted in either a syringe or I.V. bag using one of the recommended solutions. The diluted Vinorelbine Injection, USP should be administered over 6 to 10 minutes into the side port of a free-flowing I.V. followed by flushing with at least 75 to 125 mL of one of the solutions. For diluents that may be used, see below.

Syringe: The calculated dose of vinorelbine should be diluted to a concentration between 1.5 and 3 mg/mL.

I.V. Bag: The calculated dose of vinorelbine should be diluted to a concentration between 0.5 and 2 mg/mL.

Reconstituted Solutions

Syringe: Vinorelbine Injection, USP diluted to a concentration between 1.5 and 3 mg/mL may be used for up to 24 hours when stored in polypropylene syringes at 5°C to 30°C. The following solutions may be used for dilution:

- 5% Dextrose Injection, USP
- 0.9% Sodium Chloride Injection, USP

I.V. Bag: Vinorelbine Injection, USP diluted to a concentration between 0.5 and 2 mg/mL may be used for up to 24 hours when stored in polyvinylchloride bags at 5°C to 30°C. The following solutions may be used for dilution:

- 5% Dextrose Injection, USP
- 0.9% Sodium Chloride Injection, USP
- 0.45% Sodium Chloride Injection, USP
- 5% Dextrose and 0.45% Sodium Chloride Injection, USP
- Ringer's Injection, USP
- Lactated Ringer's Injection, USP

As with all the parenteral drug products, intravenous admixtures should be inspected visually for clarity, particulate matter, discoloration, and leakage prior to administration, whenever solution and container permit. Solutions showing haziness, particulate matter, precipitate, discoloration or leakage should not be used. Any unused portion should be discarded.

OVERDOSAGE

The primary anticipated complications of overdosage would consist of bone marrow suppression and peripheral neurotoxicity.

There is no known antidote for vinorelbine overdosage. Overdoses involving quantities up to ten times the recommended dose (30 mg/m²) have been reported. The toxicities described were consistent with those listed in the Adverse Reactions section including paralytic ileus, stomatitis, and esophagitis. Bone marrow aplasia, sepsis, and paresis have also been reported. Fatalities have occurred following overdose of vinorelbine tartrate. If overdosage occurs, general supportive measures together with appropriate blood transfusions, growth factors and antibiotics should be instituted as deemed necessary by the physician.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Vinorelbine tartrate is a novel vinca alkaloid which interferes with microtubule assembly. Vinca alkaloids are structurally similar compounds comprising of two multiringed units, vindoline and catharanthine. Vinorelbine is a vinca alkaloid in which the catharanthine unit is the site of structural modification. This structural change imparts unique pharmacologic properties which may translate into clinical benefits for patients with various malignancies. The antitumour activity of vinorelbine is thought to be due primarily to inhibition of mitosis at metaphase through its interaction with tubulin. Vinorelbine may also interfere with amino acid, cyclic AMP, and glutathione metabolism; calmodulin-dependent Ca⁺⁺-transport ATPase activity; cellular respiration; and nucleic acid and lipid biosynthesis.

Pharmacokinetics

Following intravenous administration, vinorelbine concentration in plasma decays in a triphasic manner. The initial rapid decline represents distribution of drug to peripheral compartments and metabolism of the drug. The prolonged terminal phase is due to relatively slow efflux of vinorelbine from peripheral compartments. The terminal phase half-life averaged 27.7 to 43.6 hours; the mean plasma clearances ranged from 0.97 to 1.26 L/hr/kg; and steady state volume of distribution (V_{ss}) values ranged from 25.4 to 40.1 L/kg.

The disposition of radiolabelled vinorelbine has been studied in a limited number of patients. Approximately 18% of the administered dose was recovered in the urine and 46% in the feces. Incomplete recovery in humans is consistent with results in animals. A separate study of the urinary excretion of vinorelbine showed that 10.9%±0.7% of a 30 mg/m² intravenous dose was excreted unchanged in the urine.

One metabolite of vinorelbine, deacetylvinorelbine, has been shown to possess antitumour activity. This metabolite has been detected but not quantified in human plasma. The effects of renal or hepatic dysfunction on the disposition of vinorelbine have not been assessed.

The pharmacokinetics of vinorelbine are not influenced by the concurrent administration of cisplatin with vinorelbine (see DRUG INTERACTIONS).

Special Populations and Conditions

Pediatrics: Pharmacokinetics of common name have not been studied in the pediatric population.

Geriatrics: Pharmacokinetics of common name have not been studied in the geriatric population.

Gender: Differences in pharmacokinetics based on gender have not been established.

Race: Differences in pharmacokinetics based on race have not been established.

Hepatic Insufficiency: Differences in pharmacokinetics based on hepatic function have not been established.

Renal Insufficiency: Differences in pharmacokinetics based on renal function have not been established.

Genetic Polymorphism: Differences in pharmacokinetics based on genetic polymorphism have not been established.

STORAGE AND STABILITY

Stability and Storage Recommendations

Store Vinorelbine Injection, USP vials refrigerated at 2°C to 8°C in the original package. Protect from light and freezing. Single use vials. Discard unused portion.

Vinorelbine Injection, USP is initially clear and colourless to pale yellow, but may develop a slightly darker yellow to light amber colour in time. This does not indicate a change which should preclude its use. Parenteral drug products should be visually inspected for particulate matter and discolouration prior to administration whenever solution and container permit. If particulate matter is seen, Vinorelbine Injection, USP should not be administered.

Syringe: Vinorelbine Injection, USP diluted to a concentration between 1.5 and 3 mg/mL may be used for up to 24 hours when stored in polypropylene syringes at 5°C to 30°C.

I.V. Bag: Vinorelbine Injection, USP diluted to a concentration between 0.5 and 2 mg/mL may be used for up to 24 hours when stored in polyvinylchloride bags at 5°C to 30°C.

SPECIAL HANDLING INSTRUCTIONS

Since vinorelbine tartrate is a cytostatic agent, established procedures specific to the handling and use of such agents must be followed.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Availability of Dosage Forms

Vinorelbine Injection, USP is supplied in clear, colourless glass vials as a solution of 10 mg / mL vinorelbine base.

Composition

Vinorelbine Injection, USP consists of an aqueous solution of vinorelbine tartrate, equivalent to 10 mg vinorelbine base per millilitre of solution. No other preservatives or other additives are present. Vinorelbine Injection, USP is a clear, colourless to pale yellow solution in Water for Injection, USP.

Supplied as:

1 mL fill in a 2 mL vial, packaged individually

5 mL fill in a 6 mL vial, packaged individually

Vial stoppers do not contain natural rubber latex.


PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance

- Proper name:** vinorelbine tartrate
Chemical name: 3',4'-didehydro-4'-deoxy-C'-norvincaleukoblastine-di-L-tartrate
Molecular formula: C₄₅H₅₄N₄O₈ • 2C₄H₆O₆
Molecular mass: Vinorelbine Tartrate: 1079.11 g/mol
Vinorelbine: 778.93 g/mol

Structural formula:



- Chemical Abstracts Service [Reg. No.]:** 125317-39-7 (Vinorelbine Ditartrate)
71486-22-1 (Vinorelbine)
105661-0702 (Vinorelbine Tartrate)

Description: Vinorelbine tartrate is a white to yellow or light brown amorphous powder having a melting point of 210 °C with decomposition. It is freely soluble in water, soluble in alcohol, slightly soluble in methylene chloride, insoluble in chloroform. pH is 3.5, pKa₁ and pKa₂ are 5.8 and 8.2, respectively.

CLINICAL TRIALS

Advanced Non-Small Cell Lung Cancer (NSCLC)

Data from two controlled clinical studies (612 + 211 patients), as well as additional data from more than 100 patients enrolled in two uncontrolled clinical trials, support the use of vinorelbine tartrate in patients with advanced non-small cell lung cancer (NSCLC). One randomized, three-arm trial in 612 Stage III or IV NSCLC patients compared treatment with single-agent

vinorelbine tartrate (30 mg/m²/week), vinorelbine tartrate (30 mg/m²/week) plus cisplatin (120 mg/m² days 1 and 29, then every six weeks), and vindesine (3 mg/m²/week for 7 weeks, then every other week) plus cisplatin (120 mg/m² days 1 and 29, then every 6 weeks). Vinorelbine tartrate, as a single agent, was well-tolerated and resulted in a median survival of 31 weeks and a response rate of 14%. The combination of vinorelbine tartrate plus cisplatin produced a median survival of 40 weeks and a response rate of 28%. The results achieved with vinorelbine tartrate as a single agent were comparable to those seen with vindesine plus cisplatin (median survival 32 weeks and response rate of 19%), but significantly less severe nausea, vomiting, and neurotoxicity were observed in patients treated with single-agent vinorelbine tartrate. In the subgroup of patients with Stage IV disease, vinorelbine tartrate plus cisplatin produced longer survival than vindesine plus cisplatin (36 weeks vs. 27 weeks).

The results with single-agent vinorelbine tartrate noted in the above trial were confirmed in a second randomized, two-arm study (211 patients) which compared treatment with single-agent vinorelbine tartrate (30 mg/m² weekly) to a control drug, 5-fluorouracil (5FU) (425 mg/m² IV bolus) plus leucovorin (LV) (20 mg/m² IV bolus) administered five consecutive days every four weeks. Compared to 5FU/LV, vinorelbine tartrate improved survival. The median survival for patients treated with vinorelbine tartrate was 30 weeks versus 22 weeks for 5FU/LV. The median survival for similar patients given best supportive care is reported to range from 9 to 21 weeks. The one year survival rates were 24% (\pm 4% S.E.) for vinorelbine tartrate and 16% (\pm 5% S.E.) for 5FU/LV, using the Kaplan-Meier product-limit estimates. The response rates for vinorelbine tartrate and 5FU/LV were 14% and 5%, respectively.

In an uncontrolled study in 78 patients with inoperable NSCLC treated with single-agent vinorelbine tartrate (30 mg/m²/week), the median survival was 33 weeks. The response rate was 33%. A Phase I/II dose-ranging study of vinorelbine tartrate (20, 25, or 30 mg/m²/week) plus cisplatin (120 mg/m² days 1 and 29, then every 6 weeks) in 32 patients with NSCLC demonstrated a median survival of 44 weeks. There were no responses at the lowest dose level; the response rate was 33% in the 21 patients treated at the two highest dose levels.

Advanced Breast Cancer

Data from one randomized, controlled clinical study (179 patients) and three uncontrolled studies (302 patients) support the effectiveness of vinorelbine tartrate in patients with advanced breast cancer. The randomized, controlled trial compared vinorelbine tartrate (30 mg/m²/week) to intravenous melphalan (25 mg/m²/every 4 weeks) in patients who had progressed on one or two prior chemotherapy regimens for advanced disease, with one regimen containing an anthracycline. Patients who relapsed during or within 6 months of treatment with an anthracycline-containing adjuvant chemotherapy regimen were also eligible. Melphalan was chosen as a Phase II comparator because of the lack of a commonly accepted standard treatment in this patient population. Treatment with vinorelbine tartrate resulted in significantly longer time to disease progression, time to treatment failure, and survival compared to melphalan. Median time to disease progression was 12 weeks for the vinorelbine tartrate patients and 8 weeks for the melphalan arm ($p < 0.001$). Median time to treatment failure was 11 weeks and 8 weeks respectively, for the vinorelbine tartrate and melphalan groups ($p < 0.001$). The median survival was 35 weeks for the group receiving vinorelbine tartrate and 31 weeks for the melphalan arm ($p = 0.03$). The 1-year survival rates were 36% and 22% respectively, for the

vinorelbine tartrate and melphalan groups. Although the proportion of patients who responded to vinorelbine tartrate (16%) was greater than those responding to melphalan (9%), the difference was not significant ($p=0.42$). However, when objective responses and stabilization were combined, the difference approached significance in favour of vinorelbine tartrate (47% vs. 28%, $p=0.06$). Disease-related symptoms improved or remained stable compared to baseline in the majority of patients in both groups.

An uncontrolled study conducted in the U.S. was designed to determine the safety and efficacy of vinorelbine tartrate (30 mg/m²/week) used as first-line or second-line therapy in the treatment of patients with advanced breast cancer. Patients were not previously treated with an anthracycline-containing regimen. The overall response rate was 35% for 60 first-line patients and 32% for 47 second-line patients. There were 9 (15%) complete responses and 12 (20%) partial responses among the first-line patients. The median duration of response was 34 weeks for both first- and second-line patients. There were 3 (6%) complete responses and 12 (26%) partial responses among second-line patients. The estimated median duration of complete response for first- and second-line patients combined exceeds 1 year.

A total of 195 patients were treated with single-agent vinorelbine tartrate (30 mg/m²/week) in two European Phase II clinical trials. The patients in both trials had no previous cytotoxic therapy for advanced breast cancer. In one trial, the objective response rate was 41% in 145 evaluable patients. In the other trial, an objective response rate of 50% was observed in 50 evaluable patients.

DETAILED PHARMACOLOGY

Pharmacodynamics

Antitumour Activity

In vitro: Vinorelbine tartrate was a potent inhibitor of tumour cell growth with broad spectrum activity comparable to vincristine (VCR) and vinblastine (VLB). Vinorelbine was shown to participate in both multi-level drug resistance (MDR) and non-MDR forms of resistance. In combination studies, an additive effect was noted when cells were exposed sequentially to vinorelbine and then to cisplatin, while synergy was observed with a simultaneous combination of vinorelbine and paclitaxel.

In vivo: Vinorelbine tartrate demonstrated antitumour activity in both murine and human tumour xenograft models that have been previously shown to be sensitive to either VLB or VCR. Vinorelbine was active by the intraperitoneal (i.p.), oral (p.o.) and intravenous (i.v.) routes on several administration schedules against i.p., i.v., and subcutaneous (s.c.) implants of various murine tumours. Vinorelbine when administered i.v. was also active against s.c. implants of lung, mammary and stomach human tumour xenografts in nude mice. Vinorelbine plus either etoposide or cisplatin provided a significant advantage in increased life span (ILS) over that of comparably dosed single agents.

Antitubulin Activity

In vitro: In cultured mouse embryos, vinorelbine tartrate was active against all classes of mitotic microtubules (the antitumour target), while it was least active against axonal microtubules (the neurotoxic target). At pharmacologic concentrations in cell-free systems, vinorelbine was much less effective than VLB or VCR at inducing spiralization of microtubules (a potential toxicity endpoint). Vinorelbine, VLB, and VCR were equipotent in inhibiting assembly regardless of isotubulins or microtubule associated proteins (MAPS).

Cell Cycle Arrest: At physiologic concentrations, vinorelbine tartrate was similar to other vinca alkaloids in its effect on tumour cell cycle kinetics, producing mitotic arrest (block in M phase without altering G2 phase) and polyploidy. Polyploid cells are usually non-viable and this probably plays a role in the mechanism of antitumour activity.

Neuropharmacological Effect in Mice and Rats: Neuropharmacologically, vinorelbine tartrate presented the same hypothermic, and acute neurotoxicity profile expected of the vinca alkaloids. Mice treated intravenously with vinorelbine experienced only slight hypothermia, which, although statistically significant, was within standard pharmacological limits. No characteristic signs that typically accompany hypothermia (e.g., ptosis, piloerection, or tremors) were noted in the vinorelbine tartrate -treated animals. Vinorelbine produced neurotoxic effects in rats after acute intravenous administration, similar to those that occurred with VCR. However, unlike VCR, vinorelbine did not produce limb paralysis. And, although mortalities occurred after both intravenous vinorelbine and VCR administration, a 10-fold greater dose of vinorelbine than VCR was required to produce the same effect.

Cardiovascular/Respiratory Effect in Dogs: Cardiovascular and respiratory studies of intravenous vinorelbine tartrate in anaesthetized dogs indicate that vinorelbine (along with two other vinca alkaloids studied) has no effect on hemodynamics, respiratory function, or the incidence of cardiac arrhythmias.

Gastrointestinal Effects in Mice and Rats: Possible gastrointestinal (GI) effects of vinorelbine tartrate were explored using models of GI transit, GI tolerance, and gastric secretion. In mice, intravenous treatment with vinorelbine had no effect on the GI transit time in a standard charcoal meal test. In mice and rats, intravenous vinorelbine showed excellent GI tolerance, with no evidence of ulcers or bleeding in the esophagus, stomach or intestine. Vinorelbine was found to inhibit gastric secretion after intravenous administration in rats. This effect, however, occurred at one half the LD₅₀ dose.

Urinary System Effects in Rats: Studies with vinorelbine tartrate on renal function indicate a moderate diuretic effect in rats after intravenous administration. Additionally, a pronounced increase in electrolyte elimination occurred. And, although there was evidence of hemolyzed blood in the urine after high intravenous doses of vinorelbine, examination of the kidneys and bladder revealed no histologic changes.

Hemobiologic Effects in Rats: Hemobiologic studies with vinorelbine tartrate showed a platelet antiaggregating property *in vitro*, but only at the very high concentration of 1 mM, which at minimum is 1000-fold the therapeutic plasma concentration.

Pharmacokinetics

In vitro studies have shown that vinorelbine is rapidly and highly distributed into cells. Studies of vinorelbine binding to human blood constituents revealed high binding to platelets and lymphocytes. The free fraction was ~ 0.11 in pooled human plasma over a concentration range of 234 to 1,169 ng/mL. The binding to plasma constituents in cancer patients ranged from 79.6% to 91.2%. Vinorelbine binding was not altered in the presence of cisplatin, 5-fluorouracil, or doxorubicin.

Absorption and Disposition: In all animal species studied (mouse, rat, dog, monkey), vinorelbine tartrate plasma concentration declined in a multi-exponential manner with an initial rapid decay followed by a slower terminal phase. Half-life values, after intravenous administration, were 7.3 hours in the mouse and 9.5 hours in the rat. Half-lives were considerably longer in the dog (ranging from 13 to 42 hours after various oral doses) and the monkey (16.6 to 35.5 hours after various oral and intravenous doses). Plasma clearances, estimated after intravenous administration in the rat and monkey, were 2 L/h/kg and 0.54 L/h/kg, respectively. Vinorelbine was shown to be rapidly absorbed after oral administration in all species studied with T_{max} values ranging from 1 to 2 hours after administration. The bioavailability of radioactivity in the mouse was 25% following a radiolabelled dose, while absorption of radioactivity was essentially complete after oral administration to the monkey. Absolute bioavailability in the rat, determined using immunoassay methodology, was found to be 16%.

Tissue Distribution: In tissue distribution studies of radiolabelled vinorelbine tartrate, radioactivity was widely distributed throughout the animals studied (mouse, rat, and monkey) with the highest amounts of radioactivity typically found in organs of elimination such as liver and kidneys. Minimal amounts of radioactivity were found in the heart and brain tissue.

Elimination and Metabolism: Vinorelbine tartrate was primarily eliminated in the feces in all species studied (mouse, rat, and monkey). Typically, 50% to 80% of an administered dose was recovered, regardless of the route of administration. A small percentage of the dose was excreted in the urine (1% to 19%, usually < 10%) and the majority of the amount excreted was recovered as unchanged drug. Evidence for substantial metabolism exists, and excretion via the bile appears to be a significant pathway for metabolites and parent drug. Three metabolites were isolated from perfused rat liver preparations, but not in sufficient quantity for identification. In vivo isolation and identification of metabolites was hindered because of the low quantities of metabolites present in animals administered non toxic doses and the inadequate sensitivity of analytical methodology.

TOXICOLOGY

Animals

Acute Toxicity

Rodents: Single-dose lethality values calculated from the results of mouse and rat studies are listed below. The combined male/female value is given, as no sex differences were apparent.

Table 4: Acute Toxicity Vinorelbine

Species/Strain	Route	LD ₁₀ (mg/kg)	LD ₅₀ (mg/kg)	LD ₉₀ (mg/kg)
Mouse/CD-1	intravenous	21.9	36.1	59.3
Mouse/OF1	intravenous	30.4	36.6	43.8
Rat/Sprague-Dawley	intravenous	10.1	11.2	12.4
Rat/Sprague-Dawley	intravenous	11.2	14.1	17.6
Mouse/CD-1	oral	56.4	82.8	121.5
Rat/Sprague-Dawley	oral	23.6	30	38.1
Mouse/CD-1	intraperitoneal	16.6	24.5	39.3
Mouse/OF1	intraperitoneal	22.7	35.5	55.5
Rat/Sprague-Dawley	intraperitoneal	6	10.1	17.1
Rat/Sprague-Dawley	intraperitoneal	2.6	6.2	14.7

Ataxia, convulsions and respiratory difficulty were noted almost immediately after dosing in animals treated intravenously at the highest doses. Deaths occurred as early as 15 minutes postdose. Clinical signs seen in animals treated by all three routes included lethargy and piloerection in mice and rats and diarrhea, chromodacryorrhea, edema of the muzzle, and prostration in rats. Swollen abdomens and evidence of peritonitis were noted in some animals treated by the intraperitoneal route. Deaths occurred from day 1 to as late as 16 and 29 days postdose in mice and rats respectively.

Lethality studies using daily doses for 5 days were conducted in mice (intravenous) and rats (oral). The 5-day cumulative mg/kg values calculated were as follows:

Table 5: Lethality Studies

Species/Strain	Route	LD ₁₀ (mg/kg)	LD ₅₀ (mg/kg)	LD ₉₀ (mg/kg)
Mouse/OF1	intravenous	31.6	39.8	49.9
Rat/Sprague-Dawley	oral	-	14	-

Mortality occurred from day 6 to 23 in mice and day 4 to 21 in rats, with clinical signs first noted on days 5 or 6. Piloerection, muzzle or facial edema, and subdued behaviour were seen in both species. In addition, skin paleness was noted in mice and diarrhea, chromorhinorrhea, and vaginal bleeding were seen in rats. Thus single and cumulative intravenous 5-dose LD₅₀s in mice were essentially the same (36 and 40 mg/kg) and the cumulative oral 5-dose LD₅₀ in rats was the same as the intravenous single-dose LD₅₀ in rats (14 and 14.1 mg/kg).

Beagle dogs: The beagle dog was the most susceptible of any species tested in single dose studies. An oral dose of 2 mg/kg and an intravenous dose of 1 mg/kg were lethal. Toxic effects included leukopenia, enteritis, elevated liver enzymes with or without histopathological evidence of hepatic necrosis or biliary hyperplasia, bone marrow hypoplasia and lymphoid depletion of spleen/lymph nodes. The lowest dose where one or more toxic effects were recorded was 0.5 mg/kg (oral and intravenous).

Monkeys: Acute toxic effects in rhesus and cynomolgus monkeys were observed after oral doses of 36 mg/kg (only dose tested) and 20 mg/kg (lowest dose) respectively and were similar to those seen in beagle dogs.

Table 6: Subchronic/Chronic Toxicity

Species/Strain	No. per Group	Route	Dose (mg/kg)	Frequency	Duration	Drug-Related Findings (Lowest Dose where Effect Noted)
Rat/ Sprague-Dawley ^a	5 M, 5 F	intravenous	1, 2, 5	1 x wk	9 wk	1 Decreased WBC/RBC, elevated liver enzymes, infections, injection site irritation 2 Decreased body weight 5 Mortality
Dog/ Beagle	2 M, 2 F	intravenous	0.25, 0.5, 0.75	1 x wk	13 wk	0.25 Injection site irritation 0.5 Decreased WBC, thymic involution, injection site irritation 0.75 Mortality, decreased RBC, increased hepatic enzymes, decreased body weight, infections
Monkey/ Rhesus	1 M, 1 F	intravenous	0.75, 1, 1.5, 2	1 x wk	29-39 wk	0.75 Decreased WBC/RBC, injection site irritation 1.5 Enteritis, decreased body weight 2 Bone marrow hypoplasia
	1 M, 1 F	intravenous	1	2 x wk	5 wk	Decreased WBC/RBC, decreased body weight, thymic involution
	1 M, 1 F	intravenous	0.2-0.8 escalating dose ^b	5 x wk, off 2 wks (1 cycle)	15 cycles	Mortality, decreased WBC/RBC, thymic involution and bone marrow hypoplasia following completion of the dosage regimen
	1 M, 1 F	intravenous	2-4 cyclic ^c	1 x wk	4 wk	Mortality, enteritis, decreased WBC/RBC, decreased body weight, bone marrow hypoplasia

Species/Strain	No. per Group	Route	Dose (mg/kg)	Frequency	Duration	Drug-Related Findings (Lowest Dose where Effect Noted)
Rat/ Sprague- Dawleya	6 M, 0 F	Oral	5, 10	2 x wk	4 wk	5 Decreased RBC 10 Mortality, decreased WBC, increased hepatic enzymes, decreased body weight, infections
	6 M, 0 F	Oral	15, 20	1 x wk	4 wk	15 Mortality, decreased RBC, increased hepatic enzymes, decreased body weight, infections 20 Decreased WBC
Rat/ Sprague- Dawleya	10 M, 10 F	Oral	1, 2, 4	daily	42 days	2 Increased hepatic enzymes 4 Mortality, decreased WBC/RBC, reduced body weight, thymic involution, bone marrow hypoplasia
	10 M, 10 F	Oral	7, 14	1 x wk	6 wk	7 Decreased WBC/RBC ^d 14 Mortality, increased hepatic enzymes, reduced body weight, thymic involution, bone marrow hypoplasia
Monkey/ Rhesus	1 M, 1 F	Oral	4-70 escalating ^e	1 x wk	13 wk	Enteritis, decreased WBC/RBC, increased hepatic enzymes, thymic involution, bone marrow hypoplasia
Monkey/ Rhesus	2 M or 2 F	Oral	4, 10	1 x wk	13 wk	4 Decreased WBC, increased hepatic enzymes
<p>^a As noted in previous studies, affected rats showed signs of piloerection, decreased activity and muzzle swelling.</p> <p>^b 0.2 mg/kg (3 cycles), 0.4 mg/kg (6 cycles), 0.6 mg/kg (3 cycles), 0.8 mg/kg (3 cycles).</p> <p>^c 4 mg/kg for 3 weeks followed by 2 mg/kg for 1 week (3 mg/kg for 4 weeks after washout in 1 animal).</p> <p>^d Splenic extramedullary hematopoiesis – 7 mg/kg once weekly dose only.</p> <p>^e 4 mg/kg escalating to 6, 8, 10, 12, 14, 20, 30, 40, 40, 50, 70, 70 (weeks 2-13).</p>						

Mutagenicity

Vinorelbine tartrate was shown to be mutagenic in a mouse micronucleus assay and an *in vivo* cytogenetics study in which bone marrow cells from treated Chinese hamsters were examined. The effect seen, polyploidy, is one that the Ames and mouse lymphoma mutagenesis assays do not detect and these assays were negative or equivocal with vinorelbine tartrate.

Reproduction and Teratology

Reproduction: No deleterious effects on maternal or fetal parameters were seen in an intravenous rat fertility/reproduction study in which males were treated once weekly for 9 weeks and females once weekly for 4 weeks with a dose of 1.5 mg/kg vinorelbine tartrate.

Teratology: Vinorelbine tartrate was found to be embryotoxic when given to mice once during gestation at 1 to 5 mg/kg intravenously but not at 0.5 mg/kg.

Other Studies

The *in vitro* hemolysis and protein flocculation test with human blood revealed a 10 mg/mL solution of vinorelbine tartrate to be unlikely to exert any hemolytic or protein flocculation problems when used clinically.

An acute study of 17-deacetyl-vinorelbine sulfate (a vinorelbine tartrate degradation product) resulted in a LD₅₀ of 23 mg/kg. Clinical signs of toxicity included subdued behaviour, prostration, piloerection and clonic convulsions. In a mouse micronucleus mutagenicity study, 17-deacetyl-vinorelbine sulfate produced an increase in the incidence in micronuclei in mice sampled 24 or 48 hours postdose. Bromo-12-vinorelbine sulfate (a vinorelbine tartrate process impurity) was negative in the same assay.

Table 7: Teratology

Species	No./ Group	Route	Dose (mg/kg) and Frequency	Drug-Related Findings
Mouse	10 F	intravenous	0.5, 1, 3, 5 One dose on day 9 of gestation	Embryotoxicity at doses 1-5 mg/kg. Fetotoxicity (delayed ossification) at 1 mg/kg.
Rat	25 F	intravenous	0.1, 0.26, 0.7 Every 3 days for 2 weeks and then through mating and until gestation day 7	Decreased weight gain and enlarged spleens in dams receiving 0.7 mg/kg. External and/or visceral abnormalities were observed in 4 fetuses at the 0.1 mg/kg dose level. Skeletal abnormalities were observed in all dose groups.
Rat	35 F	intravenous	0.1, 0.22, 0.5 Dosed on gestational days 7, 10, 13 and 16	A low incidence of skeletal abnormalities were observed in F1 fetuses in the 0.5 mg/kg dose group.
Rat	25 F	intravenous	0.1, 0.32, 1 Dosed every 3 days starting at day 17 of gestation and continuing until day 21 of lactation	Decreased weight gain was observed in female neonates during suckling and in male neonates up to 7 weeks of age at the 1 mg/kg dose only. No other deleterious effects observed.
Rabbit	6 F	intravenous	0.1, 0.25, 0.5, 0.75 Dosed on gestation days 6, 12 and 18	Decreased mean fetal weight and increased number of small fetuses, increased incidence of blood in fetal cochleas and discoloration of fetal lens observed in the 0.75 mg/kg dose group.
Rabbit	6 F	intravenous	0.1, 0.25, 0.5, 0.75 Dosed on gestation days 6, 9, 12, 15 and 18	Reduced fetal weight was observed in the 0.25 and 0.5 mg/kg dose groups. Embryotoxicity was observed at doses of 0.5 and 0.75 mg/kg. Vinorelbine tartrate was maternally lethal at 0.75 mg/kg when given for 5 but not 3 days. No terata was seen at any dose level.

Species	No./ Group	Route	Dose (mg/kg) and Frequency	Drug-Related Findings
Rabbit	15 F	intravenous	0.22, 0.4, 0.7 Dosed on gestation days 6, 9, 12, 15, 18 0.7 Dosed on gestation days 7, 10, 13, 16 0.7 Dosed on gestation days 8, 11, 14, 17	Vinorelbine tartrate was severely maternally toxic at the 0.7 mg/kg dose level and resulted in maternal clinical signs, reduced body weight and food intake, abortions, and deaths with an increased incidence of external and visceral anomalies in the few live fetuses remaining.

REFERENCES

Safe Handling of Cytotoxic Agents

1. Recommendations for the safe handling of parenteral antineoplastic drugs. Washington, DC: Division of Safety, National Institutes of Health, 1983. US Dept of Health and Human Services, Public Health Service publication NIH 83-2621.
2. AMA Council on Scientific Affairs. Guidelines for handling parenteral antineoplastics. *JAMA* 1985; 253:1590-1592.
3. National Study Commission on Cytotoxic Exposure. Recommendations for handling cytotoxic agents, 1987. Available from Louis P. Jeffrey, Chairman, National Study Commission on Cytotoxic Exposure, Massachusetts College of Pharmacy and Allied Health Sciences, 179 Longwood Avenue, Boston, Massachusetts, 02115.
4. Clinical Oncological Society of Australia. Guidelines and recommendations for safe handling of antineoplastic agents. *Med J Australia* 1983; 1:426-428.
5. Jones RB, Frank R, Mass T. Safe handling of chemotherapeutic agents: a report from the Mount Sinai Medical Center. *CA-A Cancer J for Clin* 1983; 33:258-263.
6. American Society of Hospital Pharmacists. ASHP technical assistance bulletin on handling cytotoxic and hazardous drugs. *Am J Hosp Pharm* 1990; 47:1033-1049.
7. Yodaiken RE, Bennet D. OSHA work-practice guidelines for personnel dealing with cytotoxic (antineoplastic) drugs. *Am J Hosp Pharm* 1986; 43:1193-1204.

Select Papers

8. Berthaud P, Le Chevalier T, Ruffie P, et al. Phase I-II study of vinorelbine (Navelbine[®]) plus cisplatin in advanced non-small cell lung cancer. *Eur J Cancer* 1992; 28A(11):1863-1865.
9. Binet S, Fellous A, Lataste H, et al. In-situ analysis of the action of Navelbine[®] on various types of microtubules using immunofluorescence. *Semin Oncol* 1989; 16(2, Suppl 4):5-8.
10. Cortes-Funes H, Mathé G, eds. *Drugs Supplement*. Vol 44, Suppl 4. Auckland, Adis International, 1992; 1-69.
11. Depierre A, Lemarie E, Dabouis G, et al. A phase II study of Navelbine (vinorelbine) in the treatment of non-small-cell lung cancer. *Am J Clin Oncol* 1991; 14(2):115-119.
12. Depierre A, Lemarie E, Dabouis G, et al. Efficacy of Navelbine[®] (NVB) in non-small-cell lung cancer (NSCLC). *Semin Oncol* 1989; 16(2, Suppl 4):26-29.
13. Fellous A, Ohayon R, Vacassin T, et al. Biochemical effects of Navelbine[®] on tubulin and associated proteins. *Semin Oncol* 1989; 16(2, Suppl 4):9-14.
14. Krikorian A, Rahmani R, Bromet M, et al. Pharmacokinetics and metabolism of Navelbine[®]. *Semin Oncol* 1989; 16(2, Suppl 4):21-25.

15. Pierre Fabre Oncologie (eds.) Navelbine[®] (vinorelbine) update and new trends. Paris: John Libbey Eurotext, 1991; chap. 15-22, pp.141-194.
16. Le Chevalier T, Brisgand D, Douillard JY, et al. Randomized study of vinorelbine and cisplatin versus vindesine and cisplatin versus vinorelbine alone in advanced non-small-cell lung cancer: Results of a European multicenter trial including 612 patients. *J Clin Oncol* 1994; 12(2):360-367.
17. Marty M, Extra JM, Espie M, et al. Advances in vinca-alkaloids: Navelbine[®]. *Nouv Rev Fr Hematol* 1989; 31:77-84.
18. Yokoyama A, Furusa K, Niitani H, et al. Multi-institutional phase II study of Navelbine[®] (vinorelbine) in non-small cell lung cancer, abstract No. 95. Proceedings of ASCO, March 1992; 11:287.
19. Canobbio L, Boccardo F, Pastorino G, et al. Phase II Study of Navelbine[®] in Advanced Breast Cancer, *Semin Oncol* 1989; 16(2, Suppl 4):33-36.
20. Fumoleau P, Delgado FM, Delozier T, et al. Phase II Trial of Weekly intravenous vinorelbine in first-line advanced breast cancer chemotherapy. *J Clin Oncol* 1993; 11 (7): 1245-1252.
21. Gasparini G, Caffo O, Barni S, et al. Vinorelbine is an active antiproliferative agent in pretreated advanced breast cancer patients: A Phase II study. *J Clin Oncol* 1994; 12(10):2094-2101.
22. Jones S, Winer E, Vogel C, et al. A multicentre randomized trial of IV Navelbine[®] vs. IV Alkeran[®] in patients with anthracycline-refractory advanced breast cancer (ABC). Proceedings of ASCO, March 1994; 103:22.
23. Lluch A, Garcia C, Casado A. Phase II trial with Navelbine (NVB) in advanced breast cancer (ABC) previously untreated. Proceedings of ASCO, March 1992; 11:72.
24. Marty M, Extra JM, Dieras W, et al. A review of the antitumor activity of vinorelbine in breast cancer. *Drugs* 1992; 44(Suppl 4):29-35.
25. Romero A, Rabinovich MG, Vallejo CT, et al. Vinorelbine as first-line chemotherapy for metastatic breast carcinoma. *J Clin Oncol* 1994; 12(2):336-341.
26. Spielmann M, Dorval T, Turpin F, et al. Phase II trial of vinorelbine/ doxorubicin as first-line therapy of advanced breast cancer. *J Clin Oncol* 1994; 12(9): 1764-1770.
27. Tresca P, Fumoleau P, Roche H, et al. Vinorelbine, a new active drug in breast carcinoma: Results of an ARTAC phase II trial. *Br Canc Res Treat* 1990; 16(2):161.
28. Vogel C, Bertsch L, Hohneker J. Navelbine[®]: A promising new drug with a favorable toxicity profile for breast and non-small cell lung cancer (NSCLC). *Cancer Investing* 1993; 12(Suppl 1):17-18.
29. Weber B, Vogel C, Jones S, et al. A U.S. multicenter phase II trial of Navelbine[®] in advanced breast cancer. *Proc ASCO* 1993; 12:61.
30. ONS Clinical Practice Committee. *Cancer Chemotherapy Guidelines. Recommendations for the Management of vesicant extravasation, hypersensitivity, and anaphylaxis.* Pittsburg, Pa: Oncology Nursing Society; 1992:1-4.

31. USP DI[®] System: Editorial Staff: Vinorelbine (Systemic). In: Klasco RK (Ed): USP DI[®] Advice for the Patient[®]. Thomson Micromedex, Greenwood Village, Colorado (Edition Revised 01/07/2003).
32. ^{Pr}NAVELBINE[®] (solution, 10 mg/mL), submission control 101595, Product Monograph, Pierre Fabre Pharma Canada Inc. (DEC 19, 2005)
33. ^{Pr}VINORELBINE TARTRATE FOR INJECTION (solution, 10 mg/mL), submission control 172255, Product Monograph, Teva Canada Limited. (MAR 20, 2014)

PART III: CONSUMER INFORMATION**Pr Vinorelbine Injection, USP**

This leaflet is part III of a three-part "Product Monograph" published when Vinorelbine Injection, USP was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about Vinorelbine Injection, USP. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION**What the medication is used for:**

Vinorelbine Injection, USP has been prescribed to you for the treatment of lung cancer and/or breast cancer.

What it does:

Vinorelbine interferes with the growth of rapidly dividing cells, like cancer cells, and eventually causes cell death.

When it should not be used:

Vinorelbine Injection, USP should not be used if you:

- are allergic to vinorelbine tartrate or
- who have severe granulocytopenia (reduced number of white blood cells) or severe thrombocytopenia (reduced number of platelets. Platelets help to form blood clots).

What the medicinal ingredient is:

Vinorelbine tartrate

What the important nonmedicinal ingredients are:

Vinorelbine Injection, USP: Water for Injection.

No other preservatives or other additives are present.

Vial stoppers do not contain natural rubber latex.

What dosage forms it comes in:

Solution - 10 mg vinorelbine per mL.

WARNINGS AND PRECAUTIONS**Serious Warnings and Precautions**

- **The preparation and administration of Vinorelbine Injection, USP (vinorelbine tartrate) is only carried out by trained healthcare professionals**
- **Before each administration of Vinorelbine Injection, USP, a blood sample will be taken for analysis of its components**
- **If the results of this analysis are not satisfactory, your treatment may be delayed or the dose reduced and further checks may be made**

BEFORE you use Vinorelbine Injection, USP talk to your doctor or pharmacist if:

- you have any other medical problems, especially any infections such as chickenpox (including recent exposure) or Herpes zoster (shingles);
- you are taking any medications, prescription and non-prescription medicines;
- you have had any other therapy for cancer including other drugs or radiation therapy;
- you are pregnant or if you intend to have children;
- you intend to breast-feed during treatment with vinorelbine. Vinorelbine may cause serious side effects in nursing babies;
- you have ever had any unusual or allergic reaction to vinorelbine.

INTERACTIONS WITH THIS MEDICATION

Tell your doctor about all medications that you are taking. It is especially important that your doctor knows if you are taking mitomycin, cisplatin or paclitaxel.

You may experience radiosensitivity effects if you take vinorelbine tartrate injection prior or along with radiation therapy.

PROPER USE OF THIS MEDICATION

Vinorelbine Injection, USP is for intravenous administration only. It should be administered by individuals experienced in the administration of cancer chemotherapeutic drugs.

Usual dose:

The dose of vinorelbine will be different for different patients. The dose that is used may depend on a number of things, including what the medicine is being used for, your body size, and whether or not other medicines are also being taken.

Overdose:

In the case of an overdose your doctor will stop the therapy and treat the symptoms. You may have to have additional medical treatment.

In the event of overdosage contact your doctor hospital emergency department or regional Poison Control Centre.

Missed Dose:

Your doctor will set the times at which you are to receive this medicine. It is important not to miss a dose. Let your prescriber or health care professional know if you are unable to keep an appointment. Contact your doctor as soon as possible.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Any medicine may have unwanted effects. Tell your doctor about any unusual sign or symptom whether listed or not.

Some side-effects that have been reported following treatment with vinorelbine tartrate are listed below.

- Constipation, rarely becoming intestinal obstruction
- Diarrhea
- Nausea and occasionally vomiting.
- Breathing difficulties or skin reaction as a result of an allergic reaction.
- Alopecia (hair loss).
- Jaw pain.
- A burning feeling and inflammation at the site of injection may occur. Vinorelbine tartrate is a vesicant (can cause skin blistering) and can produce phlebitis (inflammation of a vein) or extravasation (escape off vesicant drug and or blood, from a vessel into the surrounding tissue) injury. If you notice pain, swelling, burning or any unusual feeling around the site of your injection, tell your health care professional immediately.
- Decrease in white blood cells, red blood cells and platelets. The major toxicities of vinorelbine tartrate are related to bone marrow toxicity, specifically granulocytopenia (decrease in white blood cells) with

increased susceptibility to infection. Contact your doctor immediately if you experience fever or chills.

- Neuropathy (suppression of some reflex reactions or altered touch awareness).
- Asthenia (The lack or loss of strength and energy, weakness)

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
		Only if severe	In all cases	
Common	Fever, chills		√	
	Diarrhea and Constipation		√	
	Numbness of fingertips and toes		√	
	<u>At injection site</u>			
	Pain		√	
	Swelling		√	
	Burning		√	

This is not a complete list of side effects. For any unexpected effects while taking Vinorelbine Injection, USP, contact your doctor or pharmacist.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

HOW TO STORE IT

Store Vinorelbine Injection, USP vials refrigerated at 2°C to 8°C in the original package. Protect from light and freezing. Single-dose vials. Discard unused portion.

Vinorelbine Injection, USP is initially clear and colourless to pale yellow, but may develop a slightly darker yellow to light amber colour in time. This does not indicate a change which should preclude its use.

MORE INFORMATION

If you want more information about Vinorelbine Injection, USP:

- Talk to your healthcare professional.
- Find the full product monograph that is prepared for healthcare professionals by visiting the Health Canada website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website (<https://www.fresenius-kabi.com/en-ca>), or by calling 1-877-821-7724.

This leaflet was prepared by:

Fresenius Kabi Canada Ltd.
165 Galaxy Blvd, Suite 100
Toronto, ON M9W 0C8

Last Revised: August 14, 2023