


GLOBAL ANTIGEN FREQUENCIES

ABO



Antigen	ISBT	Caucasian	Black	Arabic	Indian	Chinese	Japanese	Brazilian
A	ABO1	43% ¹	27% ¹	27% ¹¹	21% ³	28% ⁴	39% ⁵	34% ^{7,8}
B	ABO2	9% ¹	20% ¹	16% ¹¹	40% ³	29% ⁴	22% ⁵	14% ^{7,8}
AB	ABO3	4% ¹	4% ¹	4% ¹¹	9% ³	9% ⁴	10% ⁵	3% ^{7,8}
O*	-	44% ¹	49% ¹	54% ¹¹	29% ³	34% ⁴	29% ⁵	49% ^{7,8}
Oh**	-	1/million ¹	-	-	1/10,000 ¹	-	1/150,000 ¹	-

Rh



Antigen	ISBT	Caucasian	Black	Arabic	Indian	Chinese	Japanese	Brazilian
D	RH1	85% ¹	92% ¹	89% ¹¹	96% ³	> 99% ²	> 99% ²	92% ^{7,8}
C	RH2	68% ¹	27% ¹	64% ¹¹	87% ⁹	92% ²	88% ²	55% ^{7,8}
CW	RH8	2% ¹	1% ¹	-	-	0% ²	0% ²	1% ^{7,8}
E	RH3	29% ¹	22% ¹	30% ¹¹	20% ⁹	40% ²	51% ²	29% ^{7,8}
c	RH4	80% ¹	96% ¹	79% ¹¹	58% ⁹	48% ²	57% ²	83% ^{7,8}
e	RH5	98% ¹	98% ¹	95% ¹¹	98% ⁹	94% ²	91% ²	92% ^{7,8}

Kell




Antigen	ISBT	Caucasian	Black	Arabic	Indian	Chinese	Japanese	Brazilian
K	KEL1	9% ¹	2% ¹	20% ¹¹	4% ⁹	< 0.01% ²	< 0.01% ²	5% ^{7,8}
k	KEL2	> 99% ¹	> 99% ¹	99% ¹¹	> 99.9% ⁹	> 99% ²	> 99% ²	95% ^{7,8}
Kpa	KEL3	2% ¹	< 0.01% ¹	-	1% ⁶	< 0.01% ²	< 0.01% ²	1% ^{7,8}
Kpb	KEL4	> 99% ¹	> 99% ¹	-	> 99% ⁶	> 99% ²	> 99% ²	> 99% ^{7,8}
Jsa	KEL6	0.01% ¹	20% ¹	-	-	-	-	4% ^{7,8}
Jsb	KEL7	> 99% ¹	99% ¹	-	-	> 99% ²	> 99% ²	> 99% ^{7,8}

Duffy




Antigen	ISBT	Caucasian	Black	Arabic	Indian	Chinese	Japanese	Brazilian
Fya	FY1	66% ¹	10% ¹	36% ¹¹	87% ⁹	> 99% ²	99% ²	39% ^{7,8}
Fyb	FY2	83% ¹	23% ¹	40% ¹¹	58% ⁹	9% ²	18% ²	50% ^{7,8}
Fy3	FY3	> 99% ¹	32% ¹	65% ¹¹	> 99% ⁶	> 99% ¹	-	70% ^{7,8}

Kidd




Antigen	ISBT	Caucasian	Black	Arabic	Indian	Chinese	Japanese	Brazilian
Jka	JK1	77% ¹	92% ¹	58% ¹¹	81% ⁹	69% ²	67% ²	93% ^{7,8}
Jkb	JK2	74% ¹	49% ¹	50% ¹¹	68% ⁹	74% ²	82% ²	54% ^{7,8}
Jk3	JK3	> 99% ¹	> 99% ¹	-	> 99% ⁹	-	-	> 99% ^{7,8}

Diego



Antigen	ISBT	Caucasian	Black	Arabic	Indian	Chinese	Japanese	Brazilian
Dia	DI1	0.01% ¹	-	-	-	4% ²	9% ²	3% ^{7,8}
Dib	DI2	> 99% ¹	> 99% ¹	-	-	-	-	> 99% ^{7,8}
Wra	DI3	0.01% ¹	0% ¹⁰	-	-	-	-	< 0.01% ^{7,8}
Wrb	DI4	> 99% ¹	-	-	-	-	-	> 99% ^{7,8}

Lewis



Antigen	ISBT	Caucasian	Black	Arabic	Indian	Chinese	Japanese	Brazilian
Lea	LE1	22% ¹	23% ¹	19% ¹¹	21% ⁶	24% ²	19% ²	19% ^{7,8}
Leb	LE2	72% ¹	55% ¹	72% ¹¹	61% ⁶	77% ²	72% ²	65% ^{7,8}

MNS



Antigen	ISBT	Caucasian	Black	Arabic	Indian	Chinese	Japanese	Brazilian
M	MNS1	78% ¹	74% ¹	96% ¹¹	89% ⁹	81% ²	79% ²	83% ^{7,8}
N	MNS2	72% ¹	75% ¹	45% ¹¹	65% ⁹	67% ²	67% ²	71% ^{7,8}
S	MNS3	55% ¹	31% ¹	65% ¹¹	55% ⁹	7% ²	10% ²	45% ^{7,8}
s	MNS4	89% ¹	93% ¹	78% ¹¹	89% ⁹	> 99% ²	> 99% ²	93% ^{7,8}
U	MNS5	> 99% ¹	99% ¹	-	> 99.9% ⁹	> 99% ²	> 99% ²	> 99% ^{7,8}
Mia***	MNS7	< 0.01% ¹²	< 0.01% ¹²	-	-	6% ¹²	< 0.01% ¹²	-

*This data refers to group O

**This data refers to Bombay which is the absence of the H antigen

*** The term 'Miltenberger' is often used to describe a complex family of hybrid glycoprotein phenotypes including GPMur, GPHut, GPHil, GPHop, etc. For more information, please refer to reference #12

References:

1. Marion Reid and Christine Lomas-Francis. The Blood Group Antigen FactsBook, Elsevier Academic Press 2005
2. M. Lin-Chu, RE Broadberry and FJ Chang. The Distribution of blood group antigens and alloantibodies among Chinese in Taiwan, Transfusion 1988; 28, 350-352
3. Tulika Chandra and Ashish Gupta. Frequency of ABO and Rhesus blood groups in blood donors, Asian Journal of Transfusion Science 2012; 6, 52-53
4. Zhao Tongmao. Genetics of Humankind blood group; Zhao Tongmao
5. Fujita Y, Tanimura, M, Tanaka K. The Distribution of the ABO Blood Groups in Japan. Jap J Human Genet 23, 63-109, 1978
6. B Thakral, K Saluja, RR Sharma, N Marwaha. Phenotype frequencies of blood groups (Rh, Kell, Kidd, Duffy, MNS, P, Lewis and Lutheran) in a North Indian blood donors, Transfusion and Apheresis Science 2010; 43, 17-22
7. Novaretti, MCZ, Dorliac-Hacer PE, Chamone DAF. Blood groups in Caucasian and Black donors from Brazil, Rev. Bras. Hematol. Hemoter. 2000, 22(1):23-32
8. Castilho L, Pellegrino Jr J. Distribution of blood groups in blood donors, Revista Paulista de Medicina 1992, 110(5): 335, 545
9. RN. Makroo, Aakanksha Bhatia, Richa Gupta & Jessy Phillip. Prevalence of Rh, Duffy, Kell, Kidd & MNSs blood group antigens in the Indian blood donor population, Indian J Med Res 137 (2013), pp 521-526
10. Issitt, P. Applied Blood Group Serology, 3rd Edition 1985 p409
11. Abdelaal MA, Anyaegbu CC, al Sobhi EM, al Baz NM, Hodan K. Blood group phenotype distribution in Saudi Arabs. Afr J Med Med Sci 1999;28:133-5.
12. Damien J. Heathcote, Timothy E. Carroll, Robert L. Flower. Sixty Years of Antibodies to MNS System Hybrid Glycoproteins: What Have We Learned? 2011; 111:124

The antigen frequencies cited in this poster have been gathered from the references listed above. If you have any studies or documentation that support antigen frequencies where data is missing, please contact your local Immucor Representative with that information for inclusion in a future revision. We appreciate your partnership!

